

UC San Diego

UC San Diego Previously Published Works

Title

A genome-wide analysis of the response to inhaled β 2-agonists in chronic obstructive pulmonary disease.

Permalink

<https://escholarship.org/uc/item/1hz62367>

Journal

The pharmacogenomics journal, 16(4)

ISSN

1470-269X

Authors

Hardin, M
Cho, MH
McDonald, M-L
et al.

Publication Date

2016-08-01

DOI

10.1038/tpj.2015.65

Peer reviewed



Published in final edited form as:

Pharmacogenomics J. 2016 August ; 16(4): 326–335. doi:10.1038/tpj.2015.65.

A Genome-wide analysis of the response to inhaled beta2-agonists in Chronic Obstructive Pulmonary Disease

Megan Hardin, MD^{1,2}, Michael H. Cho, MD^{1,2}, Merry-Lynn McDonald, PhD¹, Emily Wan, MD^{1,2}, David A. Lomas, MD³, Harvey O. Coxson, PhD⁴, William MacNee, MD⁶, Jørgen Vestbo, MD⁷, Julie C. Yates⁵, Alvar Agusti, MD⁸, Peter MA Calverley, MD⁹, Bartolome Celli, MD², Courtney Crim, MD⁵, Stephen Rennard, MD¹⁰, Emiel Wouters¹¹, Per Bakke, MD¹², Surya P Bhatt, MD¹³, Victor Kim, MD¹⁴, Joe Ramsdell, MD¹⁵, Elizabeth A. Regan, MD¹⁶, Barry J. Make, MD¹⁶, John E. Hokanson, MD¹⁷, James D. Crapo, MD¹⁶, Terri H. Beaty, PhD¹⁸, and Craig P. Hersh, MD^{1,2} on behalf of the ECLIPSE and COPD Gene Investigators

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence: Megan Hardin, MD, MPH; remeh@channing.harvard.edu, Channing Division of Network Medicine, 181 Longwood Ave, Boston, MA 02115.

Supplementary information is available at The Pharmacogenomics Journal's website.

Conflict of Interest:

The following authors report potential conflicts of interest: Dr Michael Cho receives funding from the NIH and the Alpha-1 Foundation; Dr David Lomas has received grant support, honoraria, and consultancy fees from GlaxoSmithKline. He chairs the GSK Respiratory Therapy Area Board; Dr Harvey Coxson has received \$4800 in the years 2009 – 2012 for serving on the steering committee for the ECLIPSE project for GSK, he was the co-investigator on two multi-center studies sponsored by GSK and has received travel expenses to attend meetings related to the project. Dr Coxson has three contract service agreements with GSK (including the ECLIPSE study) to quantify the CT scans in subjects with COPD and a service agreement with Spiration Inc to measure changes in lung volume in subjects with severe emphysema. He has received a fee for speaking at a conference and related travel expenses from AstraZeneca (Australia). Dr Coxson was the recipient of a GSK Clinical Scientist Award in 2010; Dr Jørgen Vestbo has received honoraria for consulting and presenting from Almirall, AstraZeneca, Boehringer-Ingelheim, Chiesi, GSK, Novartis and Takeda; Julie Yates is an employee of and owns stock in GlaxoSmithKline; Dr Alvar Agusti has consulted and received honoraria for lecturing at meeting from different pharmaceutical companies commercializing bronchodilators, including GSK, Boehringer-Ingelheim, AstraZeneca, Almirall, Novartis and Chiesi. Dr Celli has worked as a researcher or consultant for the following companies: GSK, Almirall, Novartis, Forrest, Aeris, Boehringer-Ingelheim, Dey, Altana, Pfizer, and Rox. Dr Courtney Crim is an employee of GlaxoSmithKline LLC, the sponsor of the ECLIPSE trial. He holds stock/stock options in GSK as a portion of his compensation as an employee. As it relates to this manuscript, Dr Crim declares no potential conflict of interest; Dr Rennard has had or currently has a number of relationships with companies who provide products and/or services relevant to outpatient management of chronic obstructive pulmonary disease. These relationships include serving as a consultant, advising regarding clinical trials, speaking at continuing medical education programs and performing funded research both at basic and clinical levels. He does not own any stock in any pharmaceutical companies. These companies include: AARC, American Board of Internal Medicine, Able Associates, Align2 Acton, Almirall, APT, AstraZeneca, American Thoracic Society, Beilenson, Boehringer Ingelheim, Chiesi, CIPLA, Clarus Acuity, CME Incite, COPDFoundation, Cory Paeth, CSA, CSL Behring, CTS Carmel, Dailchi Sankyo, Decision Resources, Dunn Group, Easton Associates, Elevation Pharma, FirstWord, Forest, GLG Research, Gilead, Globe Life Sciences, GlaxoSmithKline, Guidepoint, Health Advance, HealthStar, HSC Medical Education, Johnson and Johnson, Leerink Swan, LEK, McKinsey, Medical Knowledge, Medimmune, Merck, Navigant, Novartis, Nycomed, Osterman, Pearl, PeerVoice, Penn Technology, Pennside, Pfizer, Prescott, Pro Ed Communications, PriMed, Pulmatrix, Quadrant, Regeneron, Saatchi and Saatchi, Sankyo, Schering, Schlesinger Associates, Shaw Science, Strategic North, Summer Street Research, Synapse, Takeda, Telecon SC, ThinkEquity; Dr Per Bakke has consulted for Boehringer-Ingelheim and received compensation; Professor Calverley has received funding from the UK MRC and holds an NIHR programme grant. He has been compensated for work on clinical trials steering committees for GSK, Boehringer Ingelheim and Takeda. He has spoken at meetings supported by these companies and by AstraZeneca, Novartis and Almirall. He holds no stock in any relevant concern and has no contacts with the tobacco industry. Dr Victor Kim has nothing to disclose in relationship to this manuscript but has served on an advisory committee for CSA and has participated in clinical trials sponsored by Boehringer Ingelheim, Glaxo-Smith-Kline, and Roche pharmaceuticals. VK is supported by NHLBI K23HL094696-03. Dr Craig Hersh has received consulting fees from Novartis and CSL Behring.

¹Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA ²Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA ³Wolfson Institute for Biomedical Research, University College London, London, UK ⁴UBC Department of Radiology, Vancouver General Hospital, Vancouver, Canada ⁵GlaxoSmithKline, Research Triangle Park, NC, USA ⁶Centre for Inflammation Research, University of Edinburgh, Edinburgh, Scotland ⁷Department of Respiratory Medicine, Manchester Academic Health Sciences Centre, University Hospital of South Manchester, Manchester, UK ⁸Thorax Institute, Hospital Clinic, IDIBAPS, Univ Barcelona and CIBERES, SP ⁹Department of Pulmonary and Rehabilitation Medicine, University of Liverpool, Liverpool, UK ¹⁰Department of Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA ¹¹Center for Chronic Diseases, University Hospital Maastricht, Maastricht, The Netherlands ¹²Department of Clinical Science, University of Bergen, Bergen, Norway ¹³University of Alabama, Birmingham ¹⁴Division of Pulmonary and Critical Care, Temple University Hospital, Philadelphia, PA ¹⁵University of California, San Diego ¹⁶Division of Pulmonary Sciences and Critical Care Medicine, National Jewish Health, Denver, CO, USA ¹⁷Department of Epidemiology, Colorado School of Public Health, University of Colorado Denver, Denver, Colorado ¹⁸Johns Hopkins School of Public Health, Baltimore, MD, USA

Abstract

Short-acting β_2 -agonist bronchodilators are the most common medications used in treating chronic obstructive pulmonary disease (COPD). Genetic variants determining bronchodilator responsiveness (BDR) in COPD have not been identified.

We performed a genome-wide association study (GWAS) of BDR in 5789 current or former smokers with COPD in one African American and four white populations. BDR was defined as the quantitative spirometric response to inhaled β_2 -agonists. We combined results in a meta-analysis.

In the meta-analysis, SNPs in the genes *KCNK1* ($P=2.02\times 10^{-7}$) and *KCNJ2* ($P=1.79\times 10^{-7}$) were the top associations with BDR. Among African Americans, SNPs in *CDH13* were significantly associated with BDR ($P=5.1\times 10^{-9}$). A nominal association with *CDH13* was identified in a gene-based analysis in all subjects.

We identified suggestive association with BDR among COPD subjects for variants near two potassium channel genes (*KCNK1* and *KCNJ2*). SNPs in *CDH13* were significantly associated with BDR in African Americans.

Introduction

Chronic obstructive pulmonary disease (COPD) is a disorder characterized by progressive loss of lung function. It is currently the third leading cause of death world-wide, and the global burden of disease is expected to continue to rise(1). Although cigarette smoke is the greatest risk factor for COPD, recent studies have identified several genetic risk factors for this disease(2).

Inhaled bronchodilators, including β_2 -agonists, play a key role in COPD management guidelines. These medications act on smooth muscle receptors in bronchial airways to produce muscle relaxation and airway dilation, resulting in improved airflow through the lungs (1), and have been shown to alleviate COPD symptoms(3). The response to inhaled bronchodilators is measured by a change in the forced expiratory volume in one second (FEV_1) using standardized spirometry before and after the administration of β_2 -agonists. Although COPD is characterized by relatively fixed airflow limitation, up to two-thirds of COPD patients will exhibit a positive response to an inhaled bronchodilator at any one time(4).

The quantitative response to inhaled β_2 -agonists is a heritable trait(5), and candidate gene studies have identified several genes suggestive of association with quantitative measures of bronchodilator responsiveness (BDR)(6, 7). In addition, candidate gene(8) and genome-wide association studies (GWAS) have identified variants associated with BDR in subjects with asthma (9–11). We hypothesized that genome-wide association studies would identify associations with BDR in COPD.

Subjects and Methods

Study subjects

Details of the COPDGene, ECLIPSE, GenKOLS, and NETT studies, including study procedures, genotyping, and quality control, have been reported(12–16). COPDGene subjects were current and former smoking non-Hispanic white (NHW) or African American (AA) from the U.S. ECLIPSE subjects were Caucasian current or former smokers from Europe, North American and New Zealand. GenKOLS subjects were current and former smokers from Norway. NETT subjects were white former smokers from the U.S. All subjects had moderate to severe COPD (GOLD stage 2 or greater(17)). Subjects were excluded if they had a recent COPD exacerbation.

Spirometry

All subjects completed a respiratory questionnaire and performed standardized spirometry according to American Thoracic Society or European Respiratory Society guidelines. COPDGene, NETT, and GenKOLS subjects were tested before and approximately 20 minutes after administration of 2 puffs (180 μ g) of inhaled β_2 -agonist (albuterol/salbutamol). ECLIPSE subjects were tested before and 15 minutes after inhalation of 400 μ g β_2 -agonist (albuterol/salbutamol).

BDR was measured using three quantitative metrics that have been previously reported(5). BDRABS, the absolute difference in pre- versus post- bronchodilator FEV_1 ; BDRPRED, the absolute difference in pre- versus post-bronchodilator FEV_1 as a percentage of FEV_1 percent predicted; and BDRBASE, the absolute difference in pre versus post bronchodilator FEV_1 as a percentage of baseline FEV_1 .

Genotyping

All subjects were genotyped using Illumina platforms (Human Hap550 for ECLIPSE and GenKOLS, Quad610 for NETT, and Human OmniExpress for COPDGene) as previously described(13, 15). We included all variants and subjects that passed quality control, based on cluster plots (genotyped) and imputation quality ($R^2 \geq 0.80$) for imputed SNPs, Hardy-Weinberg equilibrium (P-value), and missingness (% threshold). Imputation was performed using MaCH and minimac with 1000 Genomes phase I v 3 European reference panels for white subjects. Cosmopolitan reference panels were used for COPDGene AA subjects. Variants with a minor allele frequency (MAF) $< 1\%$ and $R^2 \geq 0.80$ were excluded from analysis. Ancestry-based principal components were generated for each study using EIGENSOFT2.0(18). We performed Taqman genotyping (Applied Biosystems, Foster City, CA) for the SNPs rs114132812 and rs115067260 among 23 and 38 African American COPDGene subjects respectively, who were imputed to be carriers of the minor allele.

Statistical analysis

Baseline subject demographics and outcome variables were analyzed in R (v2.15.1). We excluded 20 subjects with BDR variables more than six standard deviations from the mean. We performed linear regression analysis for the three BDR variables in PLINK(19) including genotyped and imputed SNPs, adjusting for age, gender, pack-years smoking history, and ancestry-based principal components. We combined the results from all five samples in a fixed-effects meta-analysis using METAL(20). We additionally performed a gene-based test of significance among the top 20% of all SNPs using VEGAS (a Versatile Gene-based test for Genome-wide Association Studies). This method performs gene-based association testing by assigning SNPs within 50kb of a gene in accordance with the hg18 assembly and then uses simulation to account for linkage disequilibrium. All genes are tested for association with the trait of interest(21). Using the top 20% of significantly associated SNPs, we used VEGAS software to test 13,675 genes. Based on this number, we established a Bonferroni significance threshold of 3.6×10^{-6} . We analyzed the top genes from our GWAS as well as the top genes identified through the VEGAS analysis using the functional annotation tool, DAVID(22, 23).

We tested the association of seven asthma and three COPD SNPs previously associated with BDR in asthma GWAS (9, 11) and COPD candidate gene studies (*ADRB2*, *EPHX1*, and *SERPINE2*) (6) with BDR in our meta-analysis results. We additionally tested the association of two SNPs from the β_2 -agonist receptor gene, *ADRB2* (Arg16Gly, rs1042713 and Gln27Glu, rs1042714). We tested our top SNPs for their association with lung function (FEV₁/FVC and FEV₁) in the four COPD populations with the broadest range of lung function values: COPDGene NHW and AA, GenKOLs, ECLIPSE. We examined our top BDR variants for their association with BDR in two asthma GWAS(9, 11).

Results

The demographic data for each study population are presented in Table 1. These outcomes appear to follow a normal distribution (Supplementary Figure 1). All three outcomes are significantly correlated in COPDGene NHW, but BDRABS and BDRPRED appear more

correlated ($R^2=0.97$), compared to BDRABS and BDRBASE ($R^2=0.85$) or BDRBASE and BDRPRED ($R^2=0.88$).

The BDR outcomes were statistically different when comparing the four Caucasian populations ($P<0.05$ for ANOVA for all three BDR outcomes). However, these small differences are unlikely to be clinically significant. We additionally compared the NHW and AA subjects from COPDGene. The AA subjects demonstrated significantly lower bronchodilator responsiveness for all three outcomes, and these results remained significant, albeit small, after adjusting for age, gender, and pack-years (BDRABS: 0.10 L vs 0.07 L, $P = 0.004$; BDRPRED: 3.59% v 2.80%, $P=0.02$; BDRBASE: 9.24% vs 7.60%, $P=0.007$).

Table 2 presents the most significant SNPs from the GWAS in the non-Hispanic white subjects from COPDGene. The top SNP annotated to a gene is presented in this table. The full list of SNPs with $P < 1 \times 10^{-6}$ are presented in Supplementary Tables 1–3. SNP rs17575208, located upstream from the gene *EPHA7* on chromosome 6, was significantly associated with BDRABS ($\beta=0.11$, $P=8.29 \times 10^{-9}$). This variant was also associated with BDRPRED ($\beta=3.22$, $P=1.03 \times 10^{-7}$) and BDRBASE ($\beta=7.06$, $p=5.64 \times 10^{-6}$), though the P -values were not genome-wide significant.

Table 3 presents the top SNPs annotated to genes having $P < 5 \times 10^{-6}$ for the COPDGene AA subjects. The full list of SNPs with $P < 5 \times 10^{-6}$ are presented in Supplementary Tables 4–6. Variants in the gene *CDH13* were significantly associated with BDRABS (rs115067260; $\beta=0.17 \pm 0.03$, $P=5.05 \times 10^{-9}$) and BDRPRED (rs114132812, $\beta=7.63 \pm 1.32$, $P=1.19 \times 10^{-8}$), and showed suggestive association with BDRBASE (rs77347308 $\beta=-17.14 \pm 3.39$, $P=5.35 \times 10^{-7}$). In addition, a genotyped SNP in the gene *SGCD* was significantly associated with BDRBASE (rs10056066, $\beta=7.12 \pm 1.29$, $P=4.86 \times 10^{-8}$), and several rare imputed SNPs in the gene *GOLGA8B* were associated with the outcome BDRPRED (rs76677753, $\beta=9.49 \pm 1.67$, $P=1.9 \times 10^{-8}$). A recent GWAS using COPD subjects from the Lung Health Study population identified the variants in the gene *SGCD* as associated with airway responsiveness measured by methacholine challenge test in a physiologically distinct asthma cohort(24). While the response to inhaled methacholine is distinct from the response to inhaled bronchodilators, similar mechanisms of smooth muscle activation could be involved. We tested the two top *SGCD* SNPs from that study in our AA population. The SNP rs456290 was associated with BDRBLINE ($P=0.02$), and the SNP rs2642660 approached replication ($P=0.08$). These SNPs were not associated with BDRBLINE in the NHW population, or in the meta-analysis.

The *CDH13* SNPs were in LD (rs115067260 and rs114132812, $R^2=0.60$, $D'=1.0$). These variants were monomorphic in the Caucasian populations. We tested additional variants within *CDH13* in the COPDGene NHW subjects for association with BDR. Rs4783331 was nominally associated with BDRABS ($\beta=0.11 \pm 0.02$, $P=9.39 \times 10^{-5}$), and five additional SNPs in this gene had $P<0.001$.

We tested our model assumptions of normal distributions of the BDR traits, focusing on BDRABS in the African Americans. The BDR traits appeared to fit a normal distribution (Supplementary figure 1). We additionally examined the residuals from linear regression for

the BDR outcomes in the African American subjects, adjusted for age, pack-years, and gender, which appeared normally distributed and were consistent with our model assumption (Supplementary figure 2). We performed an inverse normal transformation of the BDRABS phenotype, and tested this trait in a GWAS. The *CDH13* SNPs, rs115067260 ($P=4.46\times 10^{-6}$) and rs114132812 ($P=5.10\times 10^{-6}$), remained the top associations. Taqman genotyping verified the imputation accuracy of the *CDH13* SNPs. 22/23 imputed carriers of the SNP rs114132812 were heterozygous for the minor allele, and for rs115067260, 35/37 subjects were verified as heterozygous and one subject homozygous for the minor allele.

We performed a meta-analysis of the results of all five study populations (Table 4, Figure 1). A SNP in the potassium channel, subfamily K, Member 1 gene (*KCNK1*) demonstrated suggestive association with the outcome BDRABS ($\beta = -0.0142$, $P=2.02\times 10^{-7}$). A SNP upstream from the gene *KCNJ2* (rs9898686) showed suggestive association with all three traits: BDRABS, ($\beta = 0.014$, $P=2.05\times 10^{-7}$), BDRBASE ($\beta = 1.26$, $P=1.83\times 10^{-7}$) and BDRPRED ($\beta = 0.44$, $P=1.22\times 10^{-6}$). Variants in the gene *MC5R* (melanocortin 5 receptor) were suggestively associated with BDRPRED (rs12956045, $\beta = -0.45$, $P=4.69\times 10^{-7}$). Several other variants in the *KCNJ2* region also demonstrated nominal association, and these variants were in linkage disequilibrium ($R^2 > 0.80$). SNPs upstream from *KCNJ2* were recently identified as associated with lung function (measured by FEV_1) in a joint meta-analysis of SNP and SNP-by-smoking effects in a population-based sample(25). We examined the top SNP from that analysis, rs11654749. This SNP was nominally associated with BDRABS in COPDGene NHW subjects, with the same effect direction ($\beta = 0.011$, $P=0.007$). We performed the meta-analysis without the AA population (Supplementary Table1), with very similar results as that including all five study populations together.

We examined SNPs previously associated with BDR in asthma GWAS (9, 11, 26, 27) (Supplementary Table 9). In asthma, the SNP rs4452682 in the gene *SLC22A23* was associated with BDR(11). This SNP was nominally associated with BDRBASE in the COPD meta-analysis ($\beta = 0.63$, $P=2.5\times 10^{-3}$), although the effect size was in the opposite direction. A rare variant in the gene *SLC24A4* was also previously associated with BDR in an asthma study(26). Although this SNP, rs77441273, was not present in our GWAS meta-analysis, several SNPs in this gene demonstrated nominal association with BDRABS (rs60243508, $\beta = 0.017$, $P=0.008$). The gene *SPATA13* was previously associated with BDR in a gene-based GWAS among African Americans with asthma(27). Although this gene did not replicate in our gene-based VEGAS analysis and the top reported SNP did not replicate in our meta-analysis, we tested all 464 of the genotyped and imputed SNPs available in the gene *SPATA13* for their association with BDRABS. Twenty-eight of these SNPs were nominally associated with BDRABS in the COPD meta-analysis, including rs9511156 ($\beta = 0.02$, $P=0.007$); however these associations were not significant after correction for multiple testing. In addition, among the COPDGene AA population, the top SNP rs9507294 from the asthma study showed nominal association with BDRBASE ($\beta = 1.94$, $P=0.05$). The remainder of the asthma GWAS SNPs were not significantly associated with BDR in the COPD analyses, and none of the asthma BDR genes were significant in the gene-based VEGAS analysis. The *ADRB2* codon 16 and 27 SNPs (rs1042713 and rs1042714) that have been previously identified in asthmatic populations were not significantly associated with BDR in the COPD GWAS. None of the top COPD variants were associated with BDR when

examined in two prior asthma GWAS(9, 11). The SNPs from candidate genes previously associated with BDR in COPD populations (6), including *EPHX1* (rs3753661), *SERPINE2* (rs6712954), and *ADRB2* (rs1042717), were not associated with BDR in our analysis.

We tested the top SNPs from the BDR meta-analysis for association with lung function in four of the five COPD populations (COPDGene NHW and AA, GenKOLs, ECLIPSE). SNP rs61824320 in the gene *KCNK1* was significantly associated with FEV₁/FVC ($\beta = -0.0042$, $P = 0.03$) and FEV₁ ($\beta = -0.02$, $P = 0.04$) (M. Cho and S. Lutz, personal communication). The remainder of top SNPs from the BDR meta-analysis were not significantly associated with lung function.

In the VEGAS gene-based analysis (Supplementary Tables 10–12), using the P value results from the meta-analysis of all five studies, *KCNK1* approached genome-wide significance for the outcome BDRABS ($P = 8.4 \times 10^{-5}$) and BDRPRED ($P = 3.8 \times 10^{-4}$). The gene SH2B adaptor protein 3, *SH2B3*, was a top gene for all three traits, approaching genome wide significance for BDRABS ($P = 3.20 \times 10^{-5}$), BDRPRED (7.0×10^{-5}) and BDRBASE (1.50×10^{-5}). We examined the top genes from the GWAS in the VEGAS analysis. *KCNJ2* and *SGCD* were not significant in the gene-based analysis; however, *CDH13* showed nominal significance for all three traits. We additionally performed Gene Ontology analysis of the top 50 genes from the GWAS and from the VEGAS analysis using the functional annotation software, DAVID. Among the top genes in the GWAS, there was enrichment for ion channel and ion transport genes, including *KCNK1*, *FXYD1*, and *PKD2L1*. Among the top genes from the VEGAS analysis, there was enrichment for chemotaxis and lipid biosynthesis.

Discussion

This is the first genome-wide association study of the response to inhaled β_2 -agonists among COPD subjects. In a meta-analysis including five COPD populations and over 5000 subjects, we identified several genetic variants associated with the response to inhaled β_2 -agonist bronchodilators. In the African-Americans from COPDGene, several variants in the genes *CDH13* (Cadherin-13), *SGCD* (Sarcoglycan delta), and *GOLGA8B* (golgin A8 family, member B) demonstrated genome-wide significance for their association with the response to β_2 -agonists. In the non-Hispanic white COPDGene population, SNPs upstream and within the potassium channel genes, *KCNJ2* and *KCNK1* respectively, approached genome-wide significant association with the response to inhaled bronchodilators. This association remained in a meta-analysis including all five COPD case populations, although not at the genome-wide significance level.

In the primary analysis among the COPDGene NHW population, the SNP rs17575208 on chromosome 6 upstream from the gene Ephrin-type A receptor 7, *EPHA7*, was genome-wide significant with a P value of 7.23×10^{-9} for the outcome BDRABS, 8.3×10^{-8} for BDRPRED and 4.15×10^{-6} for BDRBASE. Although mutations in this gene have been found in resected non-small cell lung cancer human specimens, little is known about a potential role for *EPHA7* in COPD, asthma, or BDR. In our meta-analysis, this SNP had a P value of 2.32×10^{-5} for the outcome BDRABS and 1.30×10^{-4} for the outcome BDRPRED, however the association was not significant among the other populations tested ($P > 0.05$). This is an

imputed SNP with minor allele frequency of 1.4% but good imputation quality ($R^2 = 0.82$), and therefore may suggest a promising gene for future studies.

We identified several SNPs on chromosome 17 upstream from *KCNJ2*, also known as *KIR2.1* or Potassium inwardly-rectifying channel, subfamily J, member 2, that were suggestively associated with BDR. Inwardly-rectifying potassium channels were initially described in cardiac, skeletal, and brain tissue, and especially in smooth muscle of small arterioles. These channels play a role in potassium-mediated constriction in response to hypoxemia or ischemia(28). *KCNJ2* encodes Kir2.1, a strong inwardly rectifying potassium channel, which has been identified in small and large bronchial smooth muscle cells and plays a role in membrane depolarization (29) (30). Although the exact role for these channels in smooth muscle relaxation remains to be determined, it is possible that these channels play a role in the response to increased extracellular potassium, such as that induced by hypoxemia or acidic environments, leading to membrane hyperpolarization and smooth muscle relaxation(28, 30).

Genotyped and imputed SNPs within the potassium channel gene, *KCNK1*, located on chromosome 1, were also associated with BDRABS. This gene is also known as *TWIK-1*, and encodes the 2-pore protein potassium channel subfamily K member 1, or inward rectifying potassium channel protein TWIK-1. Two-pore potassium channels have been identified in a lung epithelial cell line(31). *TWIK-1* transcript has been identified on the apical membrane of human bronchial epithelial cells, and has been suggested to play a role in hyperpolarization of membrane action potential(32). The top variant in this gene was additionally associated with measures of lung function, suggesting a potential link to COPD severity. Further studies will be necessary to confirm the roles of these potassium channel genes in BDR and COPD severity.

The identification of multiple potassium channel genes in the single SNP and gene-based analyses suggests a potential role for these channels in moderating the response to inhaled bronchodilators. Further, both the GWAS and gene-based analysis were enriched for ion channel genes. Cellular potassium levels play a key role in maintaining membrane potential, and potassium channels have been demonstrated to mediate the effects of β -agonists(2). Other potassium channel genes, such as the KCNQ voltage activated channels, have been found to ameliorate methacholine bronchoconstriction in rat lung models(33), and these inwardly-rectifying potassium channels may play a similar role in bronchial smooth muscle relaxation. The potassium channel opening medication cromakalin has been tested in animal and human asthma subjects(34). However, limited knowledge is available about the role of potassium channels in mediating smooth muscle relaxation. The identification of potassium-channel genes suggests the importance of revisiting this class of medications for COPD and asthma therapeutics.

We noted a statistically significant, but clinically small, difference in the response to inhaled β_2 -agonists between the non-Hispanic white and African American subjects in COPDGene. To our knowledge, this is the first demonstration that NHW and AA subjects with COPD may respond differently to inhaled β_2 -agonists. Because of this difference, we examined the AA population alone for variants associated with BDR. In AA subjects, SNPs in the genes

CDH13, *SGCD*, and *GOLGA8B* were associated with bronchodilator responsiveness. *CDH13* encodes the protein T-cadherin, which functions as an adiponectin receptor(35) and is expressed in mouse lungs in response to allergen stimulation with ovalbumin (36). Adiponectin is protective against allergen-induced inflammatory cell response in mouse lungs and airway hyperresponsiveness(37) and T-cadherin knock-out mice demonstrate reduced immune response, airway hyperresponsiveness, and mucus hyperplasia compared to wild-type mice (36). Elevated adiponectin levels have been associated with increased radiographic measures of percent emphysema and lower response to inhaled bronchodilators among subjects with COPD(38).

SGCD encodes the dystrophin-sarcoglycan complex protein subunit sarcoglycan- δ . This protein complex is expressed in skeletal and cardiac muscle and is thought to play a role in limb girdle muscular dystrophy(39). The delta-sarcoglycan complex has been identified in airway smooth muscle cells, and plays a role in mediating the transition of airway smooth muscle cells from contractile to proliferative phenotypes(40), suggesting a possible role in COPD pathogenesis. In addition, variants in the *SGCD* gene were recently associated with airway hyperresponsiveness in a GWAS among COPD subjects from the large multicenter Lung Health Study. These variants were also nominally associated with BDRBLINE in our population.

Variants in the gene *GOLGA8B* (Golgi A8 Family, Member B, *GOLGA8B*) were associated with BDRPRED and approached significant association with BDRABS. Although these were imputed SNPs with low minor allele frequency, the imputation was of good quality. *GOLGA8B* encodes a golgi system autoantigen, and this region has been associated with myopia in a large GWAS(41), although a potential role in bronchodilator responsiveness is unknown.

We examined the response to an inhaled bronchodilator as a quantitative variable using three measures that have been used in prior epidemiologic and genetic studies(5, 6). In a family-based study, Palmer and colleagues demonstrated that both BDRABS and BDRPRED have greater than 30% heritability, while BDRBASE is less heritable(5), suggesting that all three outcomes were suitable phenotypes to test for genetic association. The absolute change in FEV₁ after β_2 -agonist administration is the most straightforward measure, but it does not account for baseline lung function, which is reduced in COPD. In contrast, BDRBASE has been shown to correlate with baseline lung function(42). These variables were all highly correlated. As no single measure appears to be the most comprehensive, we analyzed all three traits.

Quantitative measures of the response to inhaled bronchodilators differ from the binary definition used by the American Thoracic Society and European Respiratory Society(1). Prior studies have suggested that this binary outcome does not identify a phenotypically distinct subset of COPD patients, since the presence or absence of a bronchodilator response does not predict clinical outcomes and demonstrates intra-individual variability (42, 43). In contrast, linkage and candidate gene studies have previously identified genetic risk factors for quantitative measures of bronchodilator responsiveness(5, 6). In addition to our new GWAS results, these genetic associations suggest that there are distinct genetic risk factors

that play a role in determining the quantitative response to inhaled β_2 -agonists. Identifying these markers may help to identify COPD patients who demonstrate a greater response to β_2 -agonists, or who may be unlikely to benefit and therefore should be prescribed alternative medications.

Several GWAS have identified SNPs and genes associated with BDR in asthma (8, 11, 26, 27, 44). We examined the top variants from the asthma studies for association with BDR in the COPD populations, as well as SNPs in the *ADRB2* gene that have previously been associated with clinical response to long-acting β_2 -agonist administration in COPD(44–46). A variant in the *SLC22A23* gene was nominally associated with BDR in COPD, although the effect size was in the opposite direction as originally reported. A previously identified intergenic SNP, rs11252394, on chromosome 5 also suggested significant association with BDRABS. However, none of the other asthma BDR variants demonstrated significant association. Although *ADRB2* may be a candidate gene for bronchodilator responsiveness in asthma, variants from this gene have not demonstrated consistent association with BDR in COPD (6, 47, 48).

Our study has several limitations. It is common for investigators to replicate top GWAS findings in a replication population. However, in order to improve our power to detect an association, we used all available COPD cohorts in our meta-analysis to perform the largest GWAS of BDR to date. Our most significant findings are in biologically plausible genes, and the effect sizes are similar across all included cohorts. It is encouraging that we did find some cross-over between our top hits and those in asthma populations for both bronchodilator responsiveness and airway hyperresponsiveness despite the fact that these are different study populations. Although we identified several variants upstream from *KCNJ2* as associated with BDRABS and BDRPRED, these results failed to meet genome-wide significance in the meta-analysis. An examination of each population demonstrates that these variants all had a similar effect size (Supplementary Figure 1, Supplementary Table 8), suggesting that the lack of significance may be related to sample size. In contrast, studies that have identified variants associated with lung function in the general population have had sample sizes up to ten times larger than the current study(49). This is the only genome-wide study of the response to inhaled bronchodilators in COPD performed to date. We were specifically interested in identifying genes associated with BDR in COPD populations, and therefore we are limited to available COPD cohorts for this analysis. Although the GWAS meta-analysis did not demonstrate genome-wide significance, the top SNP is upstream from the gene *KCNJ2*, and the protein-product potassium channel is relevant to the phenotype being studied. In addition, variants in this same region have previously been associated with lung function. We are additionally limited by the use of a one-time measurement of bronchodilator responsiveness. Although BDR as a binary trait is not necessarily stable in an individual COPD patient over serial measurements (43), we used quantitative outcomes in this analysis. Post-bronchodilator FEV₁ is a stable phenotype over time(5), diminishing the noise in the quantitative measures. However, the presence of intra-individual variability may have diluted our ability to identify a significant genome-wide association. Future longitudinal studies in these populations that can account for intra-individual variability may better identify genetic risk factors for these outcomes.

Within the COPDGene African American subjects, there were several SNPs that demonstrated genome-wide significance, but were of low minor allele frequency, including imputed SNPs in *CDH13*. However, all SNPs had a minor allele frequency greater than 1%, with excellent imputation quality ($R^2 = 90\%$). Although the BDR outcomes were normally distributed, minimal skewing could result in false positive associations especially among variants with low minor allele frequency. In order to test our assumption of normality, we performed inverse normal transformation of the BDR outcomes and tested for variants for association with this transformed outcome. The order of SNPs was preserved with this transformed outcome, suggesting that our assumption of normality was correct. We confirmed imputation accuracy through direct genotyping. Furthermore, animal studies provide good evidence that *CDH13* is biologically plausible as a gene potentially involved in the bronchodilator pathway. Although the top variants associated with BDR in the AA population were monomorphic in the Caucasian populations, several other *CDH13* variants were nominally associated with BDR. In addition, the gene-based test identified this gene as nominally associated with bronchodilator responsiveness even though the top SNPs in the AA analysis were not included in the gene-based test.

In summary, in the largest COPD pharmacogenetics GWAS to date, we demonstrated that variants upstream from the gene *KCNJ2* are associated with response to an inhaled short acting β_2 -agonist bronchodilator. In addition, several SNPs in the genes *CDH13* and *SGCD* were significantly associated with BDR in African Americans. These results may point to novel assessments or potential novel therapeutic pathways for COPD. Future studies will require larger COPD populations to identify genome-wide significant variants, and functional studies will help to identify a role for the SNPs and genes highlighted in the GWAS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by: U.S. National Institutes of Health grants R01 HL089897, R01 HL089856, K12 HL089990, R01 HL094635, R01 NR013377, P01 HL105339, P01 HL083069; The Sheila J. Goodnight, MD, FCCP Clinical Research Grant in Women's Lung Health; COPDGene is also supported by the COPD Foundation through contributions made to an Industry Advisory Board comprised of AstraZeneca, Pfizer, Novartis, Boehringer-Ingelheim, and Sunovion. ECLIPSE is supported by GlaxoSmithKline.

The authors would like to thank participants and field investigators in ECLIPSE, NETT, GenKOLS, and COPDGene for their willingness to contribute to medical research.

NIH Grant Support and Disclaimer

The project described was supported by Award Numbers R01 HL089897, R01 HL089856, K12 HL089990, R01 HL094635, P01 HL105339, and P01 HL083069 from the National Heart, Lung, And Blood Institute and Award Number R01 NR013377 from the National Institute of Nursing Research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Heart, Lung, And Blood Institute, the National Institute of Nursing Research, or the National Institutes of Health.

References

1. Global initiative for chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. 2013. Revised 2013 Available from: http://www.goldcopd.org/uploads/users/files/GOLD_Report_2013_Feb20.pdf
2. Foreman MG, Campos M, Celedon JC. Genes and chronic obstructive pulmonary disease. *Med Clin North Am.* 2012; 96(4):699–711. [PubMed: 22793939]
3. Ram FS, Sestini P. Regular inhaled short acting beta2 agonists for the management of stable chronic obstructive pulmonary disease: Cochrane systematic review and meta-analysis. *Thorax.* 2003; 58(7): 580–4. [PubMed: 12832670]
4. Tashkin DP, Celli B, Decramer M, Liu D, Burkhart D, Cassino C, et al. Bronchodilator responsiveness in patients with COPD. *Eur Respir J.* 2008; 31(4):742–50. [PubMed: 18256071]
5. Palmer LJ, Celedon JC, Chapman HA, Speizer FE, Weiss ST, Silverman EK. Genome-wide linkage analysis of bronchodilator responsiveness and post-bronchodilator spirometric phenotypes in chronic obstructive pulmonary disease. *Hum Mol Genet.* 2003; 12(10):1199–210. [PubMed: 12719384]
6. Kim WJ, Hersh CP, DeMeo DL, Reilly JJ, Silverman EK. Genetic association analysis of COPD candidate genes with bronchodilator responsiveness. *Respir Med.* 2009; 103(4):552–7. [PubMed: 19111454]
7. Hizawa N, Makita H, Nasuhara Y, Betsuyaku T, Itoh Y, Nagai K, et al. Beta2-adrenergic receptor genetic polymorphisms and short-term bronchodilator responses in patients with COPD. *Chest.* 2007; 132(5):1485–92. [PubMed: 17890463]
8. Duan QL, Gaume BR, Hawkins GA, Himes BE, Bleecker ER, Klanderman B, et al. Regulatory haplotypes in ARG1 are associated with altered bronchodilator response. *Am J Respir Crit Care Med.* 2011; 183(4):449–54. [PubMed: 20851928]
9. Duan QL, Lasky-Su J, Himes BE, Qiu W, Litonjua AA, Damask A, et al. A genome-wide association study of bronchodilator response in asthmatics. *Pharmacogenomics J.* 2013
10. Duan QL, Du R, Lasky-Su J, Klanderman BJ, Partch AB, Peters SP, et al. A polymorphism in the thyroid hormone receptor gene is associated with bronchodilator response in asthmatics. *Pharmacogenomics J.* 2013; 13(2):130–6. [PubMed: 22212731]
11. Himes BE, Jiang X, Hu R, Wu AC, Lasky-Su JA, Klanderman BJ, et al. Genome-wide association analysis in asthma subjects identifies SPATS2L as a novel bronchodilator response gene. *PLoS Genet.* 2012; 8(7):e1002824. [PubMed: 22792082]
12. Vestbo J, Anderson W, Coxson HO, Crim C, Dawber F, Edwards L, et al. Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE). *Eur Respir J.* 2008; 31(4): 869–73. [PubMed: 18216052]
13. Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Hersh CP, et al. Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet.* 2010; 42(3):200–2. [PubMed: 20173748]
14. Pillai SG, Ge D, Zhu G, Kong X, Shianna KV, Need AC, et al. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet.* 2009; 5(3):e1000421. [PubMed: 19300482]
15. Cho MH, Castaldi PJ, Wan ES, Siedlinski M, Hersh CP, Demeo DL, et al. A genome-wide association study of COPD identifies a susceptibility locus on chromosome 19q13. *Hum Mol Genet.* 2012; 21(4):947–57. [PubMed: 22080838]
16. Regan EA, Hokanson JE, Murphy JR, Make B, Lynch DA, Beaty TH, et al. Genetic epidemiology of COPD (COPDGene) study design. *COPD.* 2010; 7(1):32–43. [PubMed: 20214461]
17. Global Initiative for Chronic Obstructive Lung Disease. GOLD. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. Updated 2013
18. Patterson N, Price AL, Reich D. Population structure and eigenanalysis. *PLoS Genet.* 2006; 2(12):e190. [PubMed: 17194218]
19. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007; 81(3): 559–75. [PubMed: 17701901]

20. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26(17):2190–1. [PubMed: 20616382]
21. Liu JZ, McRae AF, Nyholt DR, Medland SE, Wray NR, Brown KM, et al. A versatile gene-based test for genome-wide association studies. *Am J Hum Genet*. 2010; 87(1):139–45. [PubMed: 20598278]
22. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols*. 2009; 4(1):44–57. [PubMed: 19131956]
23. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic acids research*. 2009; 37(1):1–13. [PubMed: 19033363]
24. Hansel NN, Pare PD, Rafaels N, Sin D, Sandford A, Daley D, et al. Genome Wide Association Study Identifies Novel Loci Associated with Airway Responsiveness in COPD. *Am J Respir Cell Mol Biol*. 2014
25. Hancock DB, Artigas MS, Gharib SA, Henry A, Manichaikul A, Ramasamy A, et al. Genome-wide joint meta-analysis of SNP and SNP-by-smoking interaction identifies novel loci for pulmonary function. *PLoS Genet*. 2012; 8(12):e1003098. [PubMed: 23284291]
26. Drake KA, Torgerson DG, Gignoux CR, Galanter JM, Roth LA, Huntsman S, et al. A genome-wide association study of bronchodilator response in Latinos implicates rare variants. *J Allergy Clin Immunol*. 2014; 133(2):370–8. [PubMed: 23992748]
27. Padhukasahasram B, Yang JJ, Levin AM, Yang M, Burchard EG, Kumar R, et al. Gene-based association identifies SPATA13-AS1 as a pharmacogenomic predictor of inhaled short-acting beta-agonist response in multiple population groups. *Pharmacogenomics J*. 2014; 14(4):365–71. [PubMed: 24418963]
28. Quayle JM, Nelson MT, Standen NB. ATP-sensitive and inwardly rectifying potassium channels in smooth muscle. *Physiol Rev*. 1997; 77(4):1165–232. [PubMed: 9354814]
29. Snetkov VA, Ward JP. Ion currents in smooth muscle cells from human small bronchioles: presence of an inward rectifier K⁺ current and three types of large conductance K⁺ channel. *Exp Physiol*. 1999; 84(5):835–46. [PubMed: 10502653]
30. Oonuma H, Iwasawa K, Iida H, Nagata T, Imuta H, Morita Y, et al. Inward rectifier K(+) current in human bronchial smooth muscle cells: inhibition with antisense oligonucleotides targeted to Kir2.1 mRNA. *Am J Respir Cell Mol Biol*. 2002; 26(3):371–9. [PubMed: 11867346]
31. Cook DI, Young JA. Effect of K⁺ channels in the apical plasma membrane on epithelial secretion based on secondary active Cl⁻ transport. *J Membr Biol*. 1989; 110(2):139–46. [PubMed: 2553974]
32. Zhao KQ, Xiong G, Wilber M, Cohen NA, Kreindler JL. A role for two-pore K(+) channels in modulating Na(+) absorption and Cl(-) secretion in normal human bronchial epithelial cells. *Am J Physiol Lung Cell Mol Physiol*. 2012; 302(1):L4–L12. [PubMed: 21964404]
33. Brueggemann LI, Haick JM, Neuburg S, Tate S, Randhawa D, Cribbs LL, et al. KCNQ (Kv7) POTASSIUM CHANNEL ACTIVATORS AS BRONCHODILATORS: COMBINATION WITH A beta2-ADRENERGIC AGONIST ENHANCES RELAXATION OF RAT AIRWAYS. *Am J Physiol Lung Cell Mol Physiol*. 2014
34. Pelaia G, Gallelli L, Vatrella A, Grembiale RD, Maselli R, De Sarro GB, et al. Potential role of potassium channel openers in the treatment of asthma and chronic obstructive pulmonary disease. *Life Sci*. 2002; 70(9):977–90. [PubMed: 11862989]
35. Hug C, Wang J, Ahmad NS, Bogan JS, Tsao TS, Lodish HF. T-cadherin is a receptor for hexameric and high-molecular-weight forms of Acrp30/adiponectin. *Proc Natl Acad Sci U S A*. 2004; 101(28):10308–13. [PubMed: 15210937]
36. Williams AS, Kasahara DI, Verbout NG, Fedulov AV, Zhu M, Si H, et al. Role of the adiponectin binding protein, T-cadherin (Cdh13), in allergic airways responses in mice. *PLoS One*. 2012; 7(7):e41088. [PubMed: 22815927]
37. Shore SA, Terry RD, Flynt L, Xu A, Hug C. Adiponectin attenuates allergen-induced airway inflammation and hyperresponsiveness in mice. *J Allergy Clin Immunol*. 2006; 118(2):389–95. [PubMed: 16890763]

38. Carolan BJ, Kim YI, Williams AA, Kechris K, Lutz S, Reisdorph N, et al. The association of adiponectin with computed tomography phenotypes in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2013; 188(5):561–6. [PubMed: 23777323]
39. Jung D, Duclos F, Apostol B, Straub V, Lee JC, Allamand V, et al. Characterization of delta-sarcoglycan, a novel component of the oligomeric sarcoglycan complex involved in limb-girdle muscular dystrophy. *J Biol Chem*. 1996; 271(50):32321–9. [PubMed: 8943294]
40. Sharma P, Tran T, Stelmack GL, McNeill K, Gosens R, Mutawe MM, et al. Expression of the dystrophin-glycoprotein complex is a marker for human airway smooth muscle phenotype maturation. *Am J Physiol Lung Cell Mol Physiol*. 2008; 294(1):L57–68. [PubMed: 17993586]
41. Solouki AM, Verhoeven VJ, van Duijn CM, Verkerk AJ, Ikram MK, Hysi PG, et al. A genome-wide association study identifies a susceptibility locus for refractive errors and myopia at 15q14. *Nat Genet*. 2010; 42(10):897–901. [PubMed: 20835239]
42. Calverley PM, Burge PS, Spencer S, Anderson JA, Jones PW. Bronchodilator reversibility testing in chronic obstructive pulmonary disease. *Thorax*. 2003; 58(8):659–64. [PubMed: 12885978]
43. Albert P, Agusti A, Edwards L, Tal-Singer R, Yates J, Bakke P, et al. Bronchodilator responsiveness as a phenotypic characteristic of established chronic obstructive pulmonary disease. *Thorax*. 2012; 67(8):701–8. [PubMed: 22696176]
44. Rabe KF, Fabbri LM, Israel E, Kogler H, Riemann K, Schmidt H, et al. Effect of ADRB2 polymorphisms on the efficacy of salmeterol and tiotropium in preventing COPD exacerbations: a prespecified substudy of the POET-COPD trial. *The Lancet Respiratory medicine*. 2014; 2(1):44–53. [PubMed: 24461901]
45. Chung LP, Waterer G, Thompson PJ. Pharmacogenetics of beta2 adrenergic receptor gene polymorphisms, long-acting beta-agonists and asthma. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2011; 41(3):312–26. [PubMed: 21294785]
46. Hawkins GA, Weiss ST, Bleecker ER. Clinical consequences of ADRbeta2 polymorphisms. *Pharmacogenomics*. 2008; 9(3):349–58. [PubMed: 18303970]
47. Bleecker ER, Meyers DA, Bailey WC, Sims AM, Bujac SR, Goldman M, et al. ADRB2 polymorphisms and budesonide/formoterol responses in COPD. *Chest*. 2012; 142(2):320–8. [PubMed: 22383665]
48. Hersh CP. Pharmacogenetics of chronic obstructive pulmonary disease: challenges and opportunities. *Pharmacogenomics*. 2010; 11(2):237–47. [PubMed: 20136362]
49. Hancock DB, Eijgelsheim M, Wilk JB, Gharib SA, Loefer LR, Marcianti KD, et al. Meta-analyses of genome-wide association studies identify multiple loci associated with pulmonary function. *Nat Genet*. 2010; 42(1):45–52. [PubMed: 20010835]

COPD Foundation Funding

The COPDGene® project is also supported by the COPD Foundation through contributions made to an Industry Advisory Board comprised of AstraZeneca, Boehringer Ingelheim, Novartis, Pfizer, Siemens, Sunovion, and GlaxoSmithKline

COPDGene® Investigators – Core Units

Administrative Core: James Crapo, MD (PI), Edwin Silverman, MD, PhD (PI), Barry Make, MD, Elizabeth Regan, MD, PhD

Genetic Analysis Core: Terri Beaty, PhD, Nan Laird, PhD, Christoph Lange, PhD, Michael Cho, MD, Stephanie Santorico, PhD, John Hokanson, MPH, PhD, Dawn DeMeo, MD, MPH, Nadia Hansel, MD, MPH, Craig Hersh, MD, MPH, Peter Castaldi, MD, MSc, Merry-Lynn McDonald, PhD, Emily Wan, MD, Megan Hardin, MD, Jacqueline Hetmanski, MS, Margaret Parker, MS, Marilyn Foreman, MD, Brian Hobbs, MD, Robert Busch, MD, Adel

El-Bouiez, MD, Peter Castaldi, MD, Megan Hardin, MD, Dandi Qiao, PhD, Elizabeth Regan, MD, Eitan Halper-Stromberg, Ferdouse Begum, Sungho Won, Brittney Fredericksen, Sharon Lutz, PhD

Imaging Core: David A Lynch, MB, Harvey O Coxson, PhD, MeiLan K Han, MD, MS, MD, Eric A Hoffman, PhD, Stephen Humphries MS, Francine L Jacobson, MD, Philip F Judy, PhD, Ella A Kazerooni, MD, John D Newell, Jr., MD, Elizabeth Regan, MD, James C Ross, PhD, Raul San Jose Estepar, PhD, Berend C Stoel, PhD, Juerg Tschirren, PhD, Eva van Rikxoort, PhD, Bram van Ginneken, PhD, George Washko, MD, Carla G Wilson, MS, Mustafa Al Qaisi, MD, Teresa Gray, Alex Kluiber, Tanya Mann, Jered Sieren, Douglas Stinson, Joyce Schroeder, MD, Edwin Van Beek, MD, PhD

PFT QA Core, Salt Lake City, UT: Robert Jensen, PhD

Data Coordinating Center and Biostatistics, National Jewish Health, Denver, CO: Douglas Everett, PhD, Anna Faino, MS, Matt Strand, PhD, Carla Wilson, MS *Epidemiology Core, University of Colorado Anschutz Medical Campus, Aurora, CO:* John E. Hokanson, MPH, PhD, Jennifer Black-Shinn, MPH, PhD, Gregory Kinney, MPH, PhD, Sharon Lutz, PhD, Katherine Pratte, MSPH

COPDGene® Investigators – Clinical Centers

Ann Arbor VA: Jeffrey Curtis, MD, Carlos Martinez, MD, MPH, Perry G. Pernicano, MD

Baylor College of Medicine, Houston, TX: Nicola Hanania, MD, MS, Philip Alapat, MD, Venkata Bandi, MD, Mustafa Atik, MD, Aladin Boriek, PhD, Kalpatha Guntupalli, MD, Elizabeth Guy, MD, Amit Parulekar, MD, Arun Nachiappan, MD

Brigham and Women's Hospital, Boston, MA: Dawn DeMeo, MD, MPH, Craig Hersh, MD, MPH, George Washko, MD, Francine Jacobson, MD, MPH

Columbia University, New York, NY: R. Graham Barr, MD, DrPH, Byron Thomashow, MD, John Austin, MD, Belinda D'Souza, MD, Gregory D.N. Pearson, MD, Anna Rozenshtein, MD, MPH, FACR

Duke University Medical Center, Durham, NC: Neil MacIntyre, Jr., MD, Lacey Washington, MD, H. Page McAdams, MD

Health Partners Research Foundation, Minneapolis, MN: Charlene McEvoy, MD, MPH, Joseph Tashjian, MD

Johns Hopkins University, Baltimore, MD: Robert Wise, MD, Nadia Hansel, MD, MPH, Robert Brown, MD, Karen Horton, MD, Nirupama Putcha, MD, MHS,

Los Angeles Biomedical Research Institute at Harbor UCLA Medical Center, Los Angeles, CA: Richard Casaburi, MD, Alessandra Adami, PhD, Janos Porszasz, MD, PhD, Hans Fischer, MD, PhD, Matthew Budoff, MD, Dan Cannon, PhD, Harry Rossiter, PhD

Michael E. DeBakey VAMC, Houston, TX: Amir Sharafkhaneh, MD, PhD, Charlie Lan, DO

Minneapolis VA: Christine Wendt, MD, Brian Bell, MD

Morehouse School of Medicine, Atlanta, GA: Marilyn Foreman, MD, MS, Gloria Westney, MD, MS, Eugene Berkowitz, MD, PhD

National Jewish Health, Denver, CO: Russell Bowler, MD, PhD, David Lynch, MD

Reliant Medical Group, Worcester, MA: Richard Rosiello, MD, David Pace, MD

Temple University, Philadelphia, PA: Gerard Criner, MD, David Ciccolella, MD, Francis Cordova, MD, Chandra Dass, MD, Robert D'Alonzo, DO, Parag Desai, MD, Michael Jacobs, PharmD, Steven Kelsen, MD, PhD, Victor Kim, MD, A. James Mamary, MD, Nathaniel Marchetti, DO, Aditti Satti, MD, Kartik Shenoy, MD, Robert M. Steiner, MD, Alex Swift, MD, Irene Swift, MD, Gloria Vega-Sanchez, MD

University of Alabama, Birmingham, AL: Mark Dransfield, MD, William Bailey, MD, J. Michael Wells, MD, Surya Bhatt, MD, Hrudaya Nath, MD

University of California, San Diego, CA: Joe Ramsdell, MD, Paul Friedman, MD, Xavier Soler, MD, PhD, Andrew Yen, MD

University of Iowa, Iowa City, IA: Alejandro Cornellas, MD, John Newell, Jr., MD, Brad Thompson, MD

University of Michigan, Ann Arbor, MI: MeiLan Han, MD, Ella Kazerooni, MD, Fernando Martinez, MD,

University of Minnesota, Minneapolis, MN: Joanne Billings, MD, Tadashi Allen, MD

University of Pittsburgh, Pittsburgh, PA: Frank Sciurba, MD, Divay Chandra, MD, MSc, Joel Weissfeld, MD, MPH, Carl Fuhrman, MD, Jessica Bon, MD

University of Texas Health Science Center at San Antonio, San Antonio, TX: Antonio Anzueto, MD, Sandra Adams, MD, Diego Maselli-Caceres, MD, Mario E. Ruiz, MD

ECLIPSE Principal investigators and participating centers include: Bulgaria: Y. Ivanov, Pleven; K. Kostov, Sofia. Canada: J. Bourbeau, Montreal; M. Fitzgerald, Vancouver; P. Hernández, Halifax; K. Killian, Hamilton; R. Levy, Vancouver; F. Maltais, Montreal; D. O'Donnell, Kingston. Czech Republic: J. Krepelka, Praha. Denmark: J. Vestbo, Hvidovre. The Netherlands: E. Wouters, Horn. New Zealand: D. Quinn, Wellington. Norway: P. Bakke, Bergen, Slovenia: M. Kosnik, Golnik. Spain: A. Agusti, Jaume Saulea, Palma de Mallorca. Ukraine: Y. Feschenko, Kiev; V. Gavrisyuk, Kiev; L. Yashina, Kiev. UK: L. Yashina, W. MacNee, Edinburgh; D. Singh, Manchester; J. Wedzicha, London. USA: A. Anzueto, San Antonio, TX; S. Braman, Providence, RI; R. Casaburi, Torrance, CA; B. Celli, Boston, MA; G. Giessel, Richmond, VA; M. Gotfried, Phoenix, AZ; G. Greenwald, Rancho Mirage, CA; N. Hanania, Houston, TX; D. Mahler, Lebanon, NH; B. Make, Denver, CO; S. Rennard, Omaha, NE; C. Rochester, New Haven, CT; P. Scanlon, Rochester, MN; D. Schuller,

Omaha, NE; F. Sciurba, Pittsburgh, PA; A. Sharafkhaneh, Houston, TX; T. Siler, St Charles, MO; E. Silverman, Boston, MA; A. Wanner, Miami, FL; R. Wise, Baltimore, MD; R. ZuWallack, Hartford, CT.

Steering Committee

H. Coxson (Canada), C. Crim (GlaxoSmithKline, USA), L. Edwards (GlaxoSmithKline, USA), D. Lomas (UK), W. MacNee (UK), E. Silverman (USA), R. Tal Singer (Co-chair, GlaxoSmithKline, USA), J. Vestbo (Co-chair, Denmark), J. Yates (GlaxoSmithKline, USA).

Scientific Committee

A. Agusti (Spain), P. Calverley (UK), B. Celli (USA), C. Crim (GlaxoSmithKline, USA), B. Miller (GlaxoSmithKline, USA), W. MacNee (Chair, UK), S. Rennard (USA), R. Tal-Singer (GlaxoSmithKline, USA), E. Wouters (The Netherlands), J. Yates (GlaxoSmithKline, USA).




Figure 1.
Manhattan plots for meta-analysis results for each bronchodilator responsiveness outcome




Figure 2.
Regional association plot for *KCNJ2* variants associated with the outcome BDRABS

Table 1

Study Subjects

	COPD Gene NHW	COPD Gene AA	ECLIPSE	GenKOLS	NETT
N	2792	811	1757	853	364
Population origin	US	US	Europe/US/New Zealand	Norway	US
Age	64.7 (8.2)	59.0 (8.2)	63.7 (7.1)	65.6 (10.0)	67.5 (5.8)
Males (%)	55.6	54.7	66.9	60.0	64.8
Pack-years	56.2 (27.8)	42.4 (23.1)	50.3 (27.4)	32.1 (18.5)	66.1 (30.9)
Current smoker (%)	34.7	60.8	35.4	46.9	0
BMI (kg/m ²)	28.1 (6.1)	28.0 (6.8)	26.7 (5.6)	25.3 (4.9)	25.0 (3.5)
FEV1 pp (%)	49.7 (18.0)	52.3 (17.7)	47.6 (15.6)	50.7 (17.3)	28.0 (7.4)
FEV1/FVC	0.49 (0.13)	0.53 (0.12)	0.45 (0.12)	0.51 (0.13)	0.32 (0.06)
BDRABS (L)	0.10 (0.15)	0.07 (0.18)	0.12 (0.14)	0.10 (0.14)	0.09 (0.08)
BDRBASE (%)	9.2 (12.4)	7.6 (14.2)	10.7 (13.4)	8.1 (11.7)	13.4 (12.0)
BDRPRED (%)	3.6 (4.9)	2.89 (6.5)	3.9 (4.8)	3.4 (4.7)	3.3 (3.0)

Data are presented as mean (sd) or N (%). NHW: non-Hispanic white; AA: African American; FEV1pp: FEV1 percent predicted; BMI: Body mass index. BDRABS: (Post-BD FEV1 – Pre-BD FEV1); BDRBASE: (BDRABS/(Pre-BD FEV1*100)); BDRPRED: (BDRABS/(FEV1pp*100)).

Table 2
Most significant variants and genes from the GWAS of COPD in non-Hispanic white subjects

SNP	Chr	Nearest Gene	Beta (se)	Allele	FRQ	P value
BDRABS						
rs17575208	6	<i>EPHA7</i>	0.11 (0.02)	A	0.99	8.92×10^{-9}
rs7778219 *	7	<i>LOC285889</i>	-0.03 (0.01)	G	0.11	1.62×10^{-7}
rs2367245	17	<i>KCNJ2</i>	0.02 (0.01)	G	0.57	1.33×10^{-6}
rs115310518	5	<i>CDH18</i>	0.064 (0.014)	T	0.98	2.20×10^{-6}
rs78008396	6	<i>PARK2</i>	-0.075 (0.016)	A	0.98	2.92×10^{-6}
rs7932838 *	11	<i>SOX6</i>	0.05 (0.01)	G	0.03	3.34×10^{-6}
rs11775549	8	<i>NKX2-6</i>	-0.044 (0.0095)	G	0.94	3.39×10^{-6}
rs34342951	8	<i>NKX6-3</i>	-0.021 (0.005)	G	0.72	3.42×10^{-6}
rs11260945	1	<i>IGSF21</i>	-0.046 (0.01)	G	0.95	3.74×10^{-6}
rs73671623	8	<i>STC1</i>	-0.044 (0.010)	G	0.94	4.01×10^{-6}
rs56323342	11	<i>ME3</i>	-0.040 (0.009)	G	0.95	4.21×10^{-6}
rs56010187	16	<i>HS3ST4</i>	-0.023 (0.005)	G	0.78	5.65×10^{-6}
rs1335517	14	<i>C14ORF37</i>	0.027 (0.006)	T	0.85	5.85×10^{-6}
rs181350634	3	<i>IFT80</i>	0.09 (0.02)	A	0.99	6.29×10^{-6}
rs7552783 *	1	<i>KCNK1</i>	-0.0197 (0.004)	G	0.28	7.10×10^{-6}
BDRBASE						
rs1032243	3	<i>MIR548A3</i>	-1.96 (0.40)	A	0.77	9.59×10^{-7}
rs326981	5	<i>MTRR</i>	-2.48 (0.51)	G	0.89	1.32×10^{-6}
rs4772755	13	<i>LINC00460</i>	-1.75 (0.36)	G	0.70	1.36×10^{-6}
rs6943859 *	7	<i>KLHDC10</i>	3.0 (0.62)	C	0.07	1.84×10^{-6}
rs10242432	7	<i>ZC3HC1</i>	-2.98 (0.62)	G	0.93	1.85×10^{-6}
rs8032265 *	15	<i>C15orf60</i>	3.63 (0.78)	A	0.05	3.18×10^{-6}
rs2367245	17	<i>KCNJ2</i>	1.57 (0.34)	G	0.57	3.34×10^{-6}
rs8108918 *	19	<i>VAV1</i>	1.74 (0.37)	T	0.26	3.98×10^{-6}

SNP	Chr	Nearest Gene	Beta (se)	Allele	FRQ	P value
rs1004790	2	LOC645949	-5.77(1.26)	A	0.98	4.66×10 ⁻⁶
rs17574208	6	EPHA7	7.06 (1.53)	A	0.99	5.64×10 ⁻⁶
BDRPRED						
rs17575208	6	EPHA7	3.22 (0.60)	A	0.99	1.03×10 ⁻⁷
rs7778219 *	7	LOC285889	-1.03 (0.20)	G	0.11	3.90×10 ⁻⁷
rs11775549	8	NKX2-6	-1.53 (0.31)	G	0.94	6.99×10 ⁻⁶
rs73671623	8	STC1	-1.53 (0.31)	G	0.94	1.27×10 ⁻⁶
rs2367245	17	KCNJ2	0.629 (0.13)	G	0.57	2.53×10 ⁻⁶
rs78008396	6	PARK2	-2.50 (0.53)	A	0.98	3.07×10 ⁻⁶
rs115310518	5	CDH18	2.07 (0.45)	T	0.98	4.63×10 ⁻⁶
rs1032243	3	MIR548A3	-0.73 (0.16)	A	0.77	4.30×10 ⁻⁶
rs1335517	14	C14orf37	0.89 (0.20)	T	0.85	5.65×10 ⁻⁶

Presenting the top SNP from each top gene with P < 5×10⁻⁶; Adjusted for age, sex, pack-years and principal components; Results filtered for MAF > 1%, R² > 0.80.
* Genotyped SNP.

Chr: Chromosome; FRQ: reference allele frequency; Allele: reference allele. The full list of SNPs with P < 5×10⁻⁶ is presented in the supplementary material.

Table 3
Most significant variants and genes from the GWAS of COPD in African American subjects

SNP	Chr	Nearest Gene	Beta (se)	Allele	FRQ	P value
BDRABS						
rs115067260	16	<i>CDH13</i>	0.17 (0.03)	A	0.97	5.05×10 ⁻⁹
rs140948272	17	<i>PITPNA</i>	0.33 (0.06)	C	0.99	5.24×10 ⁻⁹
rs76677753	15	<i>GOLGA8B</i>	0.25 (0.05)	A	0.99	7.96×10 ⁻⁸
rs78060357	7	<i>PLXNA4</i>	0.08 (0.02)	A	0.90	8.06×10 ⁻⁸
rs145442019	16	<i>HSBP1</i>	0.203 (0.038)	A	0.99	8.58×10 ⁻⁸
rs13345720 *	19	<i>RNA5EH2A</i>	-0.05 (0.010)	C	0.34	1.28×10 ⁻⁷
rs114871691	12	<i>BTBD11</i>	0.114 (0.02)	A	0.95	1.32×10 ⁻⁷
rs60085550	1	<i>OR6F1</i>	0.157 (0.030)	G	0.97	1.76×10 ⁻⁷
rs2295599 *	6	<i>SYCP2L</i>	0.07 (0.014)	A	0.10	5.31×10 ⁻⁷
rs149447163	2	<i>UBR3</i>	0.191 (0.038)	G	0.98	5.62×10 ⁻⁷
rs145986148	16	<i>KCNKG4</i>	0.169 (0.034)	A	0.98	6.59×10 ⁻⁷
rs12529809	6	<i>ELOVL2</i>	-0.07 (0.01)	G	0.88	8.09×10 ⁻⁷
rs387092	16	<i>MLYCD</i>	0.20 (0.04)	A	0.99	9.90×10 ⁻⁷
rs4742936	9	<i>ABCA1</i>	0.088 (0.018)	C	0.94	9.95×10 ⁻⁷
BDRBASE						
rs10056066 *	5	<i>SGCD</i> *	7.12 (1.29)	A	0.07	4.86×10 ⁻⁸
rs143376495	6	<i>TFAP2B</i>	-19.47 (3.63)	G	0.99	1.06×10 ⁻⁷
rs2295599 *	6	<i>SYCP2L</i> *	6.0 (1.13)	A	0.10	1.38×10 ⁻⁷
rs11651753 *	17	<i>MIR4315-2</i>	4.22 (0.815)	T	0.22	2.81×10 ⁻⁷
rs75762663	13	<i>PRR20E</i>	-19.22 (3.80)	A	0.99	5.34×10 ⁻⁷
rs77347308	16	<i>CDH13</i>	-17.14 (3.39)	A	0.99	5.35×10 ⁻⁷
rs77094751	18	<i>SALL3</i>	-10.23 (2.03)	G	0.96	5.65×10 ⁻⁷
rs62385074	5	<i>LCP2</i>	-10.17 (2.03)	G	0.96	6.32×10 ⁻⁷
rs141697850	4	<i>TRAM1L1</i>	-15.05 (3.01)	C	0.99	7.02×10 ⁻⁷

SNP	Chr	Nearest Gene	Beta (se)	Allele	FRQ	P value
rs12529809	6	ELOVL2	-5.674 (1.14)	G	0.88	7.85×10 ⁻⁷
rs2575515	4	GRXCRI	3.525 (0.71)	T	0.47	8.33×10 ⁻⁷
rs7848679	9	ABCA1	6.96 (1.41)	G	0.93	9.31×10 ⁻⁷
rs76999017	2	INPP5D	-8.263 (1.67)	G	0.95	9.70×10 ⁻⁷
BDRPRED						
rs114132812	16	CDHL3	7.63 (1.32)	A	0.98	1.19×10 ⁻⁸
rs76677753	15	GOLGA8B	9.49 (1.67)	A	0.99	1.90×10 ⁻⁸
rs13345720*	19	RNASEH2A	-1.83 (0.35)	C	0.34	1.60×10 ⁻⁷
rs114871691	12	BTBD11	4.116 (0.79)	A	0.95	2.06×10 ⁻⁷
rs72653721	6	SYCP2L	-2.746 (0.53)	C	0.88	2.91×10 ⁻⁷
rs147388556	1	LPHN2	4.17 (0.81)	T	0.96	3.32×10 ⁻⁷
rs11245387	10	FAM175B	-1.76 (0.34)	T	0.40	3.49×10 ⁻⁷
rs145442019	16	HSBP1	7.10 (1.39)	A	0.99	3.55×10 ⁻⁷
rs143859190	2	GALNT13	8.10 (1.58)	T	0.99	3.67×10 ⁻⁷
rs188637550	8	MTUS	6.89 (1.36)	T	0.98	4.60×10 ⁻⁷
rs78060357	7	PLXNA4	2.88 (0.57)	A	0.90	5.08×10 ⁻⁷
rs12529809	6	ELOVL2	-2.64 (0.52)	G	0.88	5.37×10 ⁻⁷
rs188524320	8	PCMI	7.01 (1.40)	C	0.99	6.39×10 ⁻⁷
rs4742936	9	ABCA1	3.28 (0.66)	C	0.94	7.17×10 ⁻⁷
rs17116498	8	FGL1	7.07 (1.42)	C	0.99	7.32×10 ⁻⁷
rs139218005	8	ATP6V1H	4.935 (0.89)	A	0.97	9.74×10 ⁻⁷

Presenting the top SNP from each top gene with $P < 1 \times 10^{-6}$; Adjusted for age, sex, pack-years and principal components; Results filtered for MAF $> 1\%$, $R^2 > 0.80$.
* Genotyped SNP.

Chr: Chromosome; FRQ: reference allele frequency; Allele: reference allele. The full list of SNPs with $P < 5 \times 10^{-6}$ is presented in the supplementary material.

Table 4

Meta-analysis of top variants associated with BDR from 5 COPD populations

SNP	Chr	Gene	Allele	Effect	Std Err	P Value
BDRABS						
rs61824320	1	KCNK1	A	-0.0142	0.0027	2.02×10 ⁻⁷
rs9898686	17	KCNJ2	T	0.0138	0.0027	2.05×10 ⁻⁷
rs68120025	17	KCNJ2	C	0.0138	0.0027	2.05×10 ⁻⁷
rs68193035	17	KCNJ2	A	0.0138	0.0027	2.05×10 ⁻⁷
rs9906150	17	KCNJ2	A	0.0137	0.0026	2.06×10 ⁻⁷
rs9899756	17	KCNJ2	C	0.0138	0.0027	2.08×10 ⁻⁷
rs67158616	17	KCNJ2	C	0.014	0.0028	5.35×10 ⁻⁷
rs12956045	18	MC5R	A	-0.013	0.0026	8.30×10 ⁻⁷
BDRPRED						
rs68193035	17	KCNJ2	A	1.26	0.24	1.79×10 ⁻⁷
rs68120025	17	KCNJ2	C	1.26	0.24	1.79×10 ⁻⁷
rs9906150	17	KCNJ2	A	1.25	0.24	1.82×10 ⁻⁷
rs9898686	17	KCNJ2	T	1.26	0.24	1.83×10 ⁻⁷
rs9899756	17	KCNJ2	C	1.25	0.24	1.85×10 ⁻⁷
rs11871999	17	KCNJ2	T	1.26	0.26	9.36×10 ⁻⁷
BDRBASE						
rs12956045	18	MC5R	A	-0.45	0.09	4.69×10 ⁻⁷
rs2278843	10	PKD2LI	G	0.42	0.08	7.11×10 ⁻⁷
rs7079679	10	PKD2LI	A	0.42	0.08	7.26×10 ⁻⁷
rs28650403	19	NWDI	G	-1.26	0.26	9.82×10 ⁻⁷
rs28428860	19	NWDI	C	-1.26	0.26	9.94×10 ⁻⁷
rs28668077	19	NWDI	C	-1.26	0.26	9.95×10 ⁻⁷

Most significant variants associated with BDR among the meta-analysis including COPDGene NHW and AA, ECLIPSE, GenKOLS, and NETT. Allele: reference allele.